



**UNIVERSITI PUTRA MALAYSIA**

**POTASSIUM DYNAMIC AND AVAILABILITY  
FROM COMPOSTED AND UNCOMPOSTED  
RICE STRAW**

**MULYADI**

**FP 2000 22**

**POTASSIUM DYNAMIC AND AVAILABILITY  
FROM COMPOSTED AND UNCOMPOSTED  
RICE STRAW**

**MULYADI**

**MASTER OF AGRICULTURAL SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**2000**

**POTASSIUM DYNAMIC AND AVAILABILITY  
FROM COMPOSTED AND UNCOMPOSTED RICE STRAW**

**By**

**MULYADI**

**Thesis Submitted in Fulfilment of the Requirement for the  
Degree of Master of Agricultural Science in the Faculty of Agriculture  
Universiti Putra Malaysia**

**December 2000**



*Dedicated to*  
*Allah S. W. T.,*  
*my Parents,*  
*my wife Dyah Tri M. H.,*  
*my children Alfistia Md.*  
*and Yogi C. A.*

Abstract of the thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirement of the degree of Master of Agricultural Science.

**POTASSIUM DYNAMIC AND AVAILABILITY  
FROM COMPOSTED AND UNCOMPOSTED RICE STRAW**

**By**

**MULYADI**

**December 2000**

**Chairman: Ahmad Husni Mohd. Hanif, Ph.D.**

**Faculty: Agriculture**

Managing rice straw (in term of K source) is essential if sustainability is to be achieved with a small amount of inorganic K fertilizer. About 80 % K taken up by rice crop (*Oryza sativa* L.) is in the straw. To increase efficiency of K utilization from the rice straw, it is essential to know the K supplying characteristic and crop available K from the straw. Information of this kind may also help in a better understanding of K cycling in soil-crop system.

Two experiments were conducted in a glasshouse. The first experiment, composting of rice straw, was to study the changes in tensile load of rice straw during rice straw composting, its possibility as an indicator for the state of decomposition and K released. The second experiment, a pot experiment using corn (*Zea mays* L.) variety of PJ-58 as test crop for K uptake planted on Bungor series soil (Typic Paleudult), was firstly to evaluate the crop available K from composted (CRS) and uncomposted (UCRS) rice straw compared to that from Muriate of Potash (MOP) as standard K



fertilizer and secondly to evaluate the use of tensile load of rice straw as an indicator to predict K released from UCRS incorporated in the soil, the crop K uptake and leaf K concentration.

The results indicated that during composting of rice straw, the individual relationships between the percentage of organic matter remaining, the contents of total and water soluble K of decomposing rice straw with the tensile load of indicator rice straw are highly significant ( $P \leq 0.01$ ) with linear correlation coefficients of 0.97, - 0.96 and - 0.94, respectively. Therefore, tensile load of indicator rice straw can possibly be used as an alternative indicator for the state of decomposition of rice straw and to predict K released. Applications of MOP, CRS and UCRS to the soil increased K accumulated in corn crop, total and exchangeable K contents of the soil, but their increase depends on the crop growth period, the K rate and availability of fertilizer applied. Compared to MOP and UCRS, the use of CRS is more beneficial in increasing the crop K, P and Ca uptake and results in better crop growth. As a source of K, the K availability from CRS was more readily available than that from UCRS and MOP; however at 56 days after planting (tasseling stage) and a rate of 360 mg K pot<sup>-1</sup> (90 kg K ha<sup>-1</sup>), the crop available (uptake) K from the three K fertilizer sources were similar, ranging from 90.48 to 109.25 %. Tensile load of indicator rice straw can also possibly be used as indicator to predict K released from UCRS incorporated into the soil based on the correlation of tensile load and K content in the decomposing rice straw during composting. The trends are less applicable as indicator to predict the crop K uptake and leaf K concentration of corn.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains Pertanian.

**KEDINAMIKAN DAN KEDAPATAN KALIUM DARIPADA  
JERAMI PADI YANG DIKOMPOS DAN TIDAK DIKOMPOS**

**Oleh**

**MULYADI**

**Disember 2000**

**Pengerusi: Ahmad Husni Mohd. Hanif, Ph.D.**

**Fakulti: Pertanian**

Pengurusan jerami padi (dalam konteks sumber K) adalah penting untuk mencapai kelestarian dengan pertambahan baja K tak organik yang sedikit. Lebih kurang 80 % K yang diserap oleh padi (*Oryza sativa* L.) adalah di dalam jeraminya. Untuk meningkatkan kecekapan penggunaan K daripada jerami padi, adalah penting untuk mengetahui sifat pembekalan K serta K tersedia untuk tanaman daripada jerami padi. Maklumat sebegini membantu pemahaman yang lebih mendalam mengenai kitaran K di dalam sistem tanah-tanaman.

Dua percubaan telah dijalankan di rumah kaca. Percubaan pertama, pengkomposan jerami padi, adalah untuk melihat perubahan galasan tensil jerami padi semasa proses pengkomposan jerami padi, dan kemungkinan ia sebagai penunjuk untuk menyatakan pereputan dan K yang dibebaskan. Percubaan kedua, menggunakan tanaman jagung (*Zea mays* L.) varieti PJ-58 sebagai tanaman ujian untuk

pengambilan K yang ditanam pada tanah siri Bungor (Tipik Paleudult), adalah pertamanya untuk menilai K tersedia untuk tanaman daripada jerami padi dikompos (CRS) dan jerami padi tidak dikompos (UCRS) berbanding dengan “Muriate of Potash” (MOP) sebagai baja K yang piawai dan keduanya untuk menilai kegunaan galasan tensil jerami padi sebagai penunjuk untuk meramal K yang dibebaskan daripada jerami padi yang dimasukkan ke dalam tanah, pengambilan K oleh tanaman dan kandungan K daun.

Keputusan-keputusan menunjukkan bahawa semasa pengkomposan jerami padi, hubungan antara peratus sisa bahan organik, kandungan jumlah K dan kandungan K larut air daripada jerami padi yang sedang mereput dengan galasan tensil daripada jerami penanda adalah sangat nyata ( $P \leq 0.01$ ) dengan koefisien korelasi linear masing-masing 0.97, - 0.96 and - 0.94. Oleh itu, galasan tensil daripada jerami penanda boleh diguna sebagai penunjuk alternatif untuk menyatakan pereputan jerami padi dan untuk meramal K yang dibebaskan. Pemberian daripada MOP, CRS and UCRS ke tanah meningkatkan K yang dikumpul dalam tanaman jagung, kandungan jumlah K dan K tukarganti dalam tanah, tetapi peningkatan ini bergantung pada jangka waktu tumbesaran tanaman, kadar dan kedapatan K daripada baja yang diguna. Dibanding dengan MOP dan UCRS, penggunaan CRS adalah lebih baik dalam peningkatan pengambilan K, P dan Ca oleh tanaman dan menghasilkan tumbesaran tanaman yang lebih baik. Sebagai sumber K, kedapatan K daripada CRS adalah lebih cepat tersedia berbanding kedapatan K daripada UCRS dan MOP, tetapi pada 56 hari selepas ditanam (tahap pembungaan) dan pada kadar 360 mg K per pasu ( $90 \text{ kg K ha}^{-1}$ ), K tersedia untuk tanaman daripada tiga



sumber baja K itu tidak berbeza, ia berkisar dari 90.48 ke 109.25 %. Galasan tensil daripada jerami penanda mungkin juga diguna sebagai penunjuk untuk meramal K yang dibebaskan daripada UCRS yang dimasukkan ke dalam tanah berdasarkan korelasi daripada galasan tensil dan kandungan K dalam jerami padi yang mereput selama pengkomposan. Tren galasan tensil kurang berguna sebagai penunjuk untuk meramal pengambilan K dan kandungan K dalam daun jagung.

## ACKNOWLEDGEMENTS

The author wishes to express this sincere gratitude to Dr. Ahmad Husni Mohd. Hanif the, Chairman of the Supervisory Committee, for his keen interest, valuable contribution and tiredness guidance during the preparation of this thesis. His countless patience, encouragement and generosity cannot be over emphasized.

The author is also very grateful to Dr. Anuar Abdul Rahim, Dr. Halimi Mohd. Saud and Dr. Azmi Yahya, the members of the Advisory Committee, for their invaluable assistance and guidance at the various stages for my research and preparing of this thesis.

The support of the Government of Indonesia through the Agency for Agricultural Research and Development, particularly to Agricultural Research Management Project phase II for providing financial assistance is gratefully acknowledged.

I am greatly indebted to the entire technical staff of the Land Management Department especially to Puan Sarimah Hashim, Mr. Abdul Rahim Utar, Mr. Jamil Omar, Mr. Mohd Fuzi Mohd Sharif and Mr. Junaidi Jaafar as well as to Mr Mohd. Roshdi Zamri, technical staff of Biological and Agricultural Engineering, for their diverse cooperation that led to the smooth running of experiments.

I certify that an Examination Committee met on 1<sup>st</sup> December 2000 to conduct the final examination of Mulyadi on his Master of Agricultural Science thesis entitled “Potassium Dynamic and Availability from Composted and Uncomposted Rice Straw” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Mohd. Khanif Yusop, Ph.D.**

Associate Professor,  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Ahmad Husni Mohd. Hanif, Ph.D.**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Anuar Abdul Rahim, Ph.D.**


Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Halimi Mohd. Saud, Ph.D.**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Azmi Yahya, Ph.D.**

Faculty of Engineering  
Universiti Putra Malaysia  
(Member)



---

**MOHD. GHAZALI MOHAYIDIN, Ph.D.**  
Professor/Deputy Dean of Graduate School,  
Universiti Putra Malaysia

Date: **12 DEC 2000**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science.



KAMIS AWANG, Ph.D.  
Associate Professor  
Dean of Graduate School,  
Universiti Putra Malaysia

Date: 11 JAN 2001

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



**MULYADI**

Date: 11-12-2000

## TABLE OF CONTENTS

	Page
<b>DEDICATION.....</b>	ii
<b>ABSTRACT.....</b>	iii
<b>ABSTRAK.....</b>	v
<b>ACKNOWLEDGEMENTS.....</b>	viii
<b>APPROVAL SHEETS.....</b>	ix
<b>DECLARATION FORM.....</b>	xi
<b>LIST OF TABLES.....</b>	xv
<b>LIST OF FIGURES.....</b>	xvi
<b>LIST OF ABBREVIATIONS.....</b>	xviii
<b>CHAPTER</b>	
<b>I INTRODUCTION.....</b>	1
<b>II LITERATURE REVIEW.....</b>	4
Potassium Requirement and the Function in Crop Growth and Yield	4
Potassium in Soil.....	5
Bearing Minerals and Other Sources of K in Soil.....	6
Forms and Availability of Soil Potassium.....	7
Potassium Dynamics in Soil-Plant System.....	8
Potassium and Factors Influencing Uptake .....	10
Potassium Fertilizers and Organic Manure as Source of Nutrients...	11
Potential of Rice Straw as a Source of Potassium, and its Effect on Soil and Crop Yield.....	12
Organic Material Decomposition, Factors Influencing and Nutrient Release .....	14
The Role of Microorganisms in Decomposition of Organic Material.....	15
Composting.....	18
Decomposition of Organic Materials Applied to Soil.....	20
Weight Loss and Tensile Load as Indicator for Decomposition of Organic Materials Applied to Soil.....	23
<b>III POTASSIUM AND RICE STRAW TENSILE LOAD CHANGES DURING COMPOSTING OF RICE STRAW.....</b>	24
Introduction.....	24
Materials and Methods.....	25
Materials.....	25
Method of Composting.....	28
Sampling and Measurement of Tensile Load of Rice Straw	29
Sampling and Preparation of Composted Rice Straw for Chemical Analyses.....	29
Chemical Analyses of Composted Rice Straw.....	30

Data Analysis.....	32
Results and Discussion .....	33
Temperature Variations during Composting Process.....	33
Organic Matter Decomposition and Stabilization during Composting.....	34
Changes in Tensile Load of Decomposing Indicator Rice Straw, and the Relationship with Organic Matter Decomposition during Composting.....	36
Changes in Potassium Content, and the Relationship with Tensile Load of Indicator Rice Straw during Composting...	38
Conclusions.....	40
 <b>IV POTASSIUM AVAILABILITY FROM COMPOSTED AND UNCOMPOSTED RICE STRAW.....</b>	 41
Introduction.....	41
Materials and Methods.....	42
Site of Experiment and Materials.....	42
Experimental Design and Treatments.....	45
Estimation of Rice Straw Decomposition and K Released	46
Soil Sampling and Data Collected After Planting of Corn...	47
Corn Plant Sampling and Related Data .....	48
Physico-Chemical Analyses of Soil.....	49
Chemical Analyses of Urea, TSP, MOP and GML.....	51
Chemical Analyses of Composted and Uncomposted Rice Straw.....	52
Chemical Analyses of Corn Plant.....	54
Data Analysis.....	55
Results and Discussion .....	55
Effects of K Applied from MOP, Composted and Uncomposted Rice Straw on Corn Crop Growth.....	55
Effects of K Applied from MOP, Composted and Uncomposted Rice Straw on Crop K, P, Ca and Mg Uptake, and the Apparent K-fertilizer Recovery.....	57
Effects of K Applied from MOP, Composted and Uncomposted Rice Straw on Soil Total and Exchangeable K Contents.....	70
Changes in Tensile Load of Decomposing Indicator Rice Straw in Soil, and the Relationship with Crop K Uptake and Leaf K Concentration of Corn .....	77
Conclusions.....	82
 <b>V SUMMARY OF DISCUSSION.....</b>	 84
 <b>VI CONCLUSIONS.....</b>	 91
 <b>REFERENCES.....</b>	 93

<b>APPENDIX.....</b>	<b>100</b>
<b>BIODATA OF AUTHOR.....</b>	<b>103</b>



## LIST OF TABLES

		<b>Page</b>
<b>Table</b>		
1	Chemical Properties of Raw Materials Used in Composting.....	26
2	Initial Physico-Chemical Characteristics of Soil Used in Experiment..	43
3	The Contents of Organic Matter and Nutrient Elements of Fertilizers Used in Experiment.....	44
4	Treatments in Pot Experiment Using Corn .....	45
5	Mean Comparison for Dry Matter Yield and Crop Nutrient (K, P, Ca and Mg) Uptake of Corn on Soil Treated with Different Rates and Sources of K at Different Growth Periods. The Comparison using Dunnett Test at Significant Level of 5 % and the Standard Treatment is 360 mg K pot <sup>-1</sup> from Muriate of Potash (MOP).....	100
6	Mean Comparison for Apparent K Recovery from Different Rates and Sources of K at Different Growth Periods. The Comparison using Dunnett Test at Significant Level of 5 % and the Standard Treatment is 360 mg K pot <sup>-1</sup> from Muriate of Potash (MOP).....	101
7	Mean Comparison for Total and Exchangeable K Contents of Soil Treated with Different Rates and Sources of K Fertilizer at Different Growth Periods of Corn Crop. The Comparison using Dunnett Test at Significant Level of 5 % and the Standard Treatment is 360 mg K pot <sup>-1</sup> from Muriate of Potash (MOP).....	102
8	Mean Comparison for Tensile Load of Decomposed Indicator Rice Straw during 7, 14, 28 and 56 days in Soil treated with Uncomposted Rice Straw (UCRS) Equivalent to 180, 360 and 540 mg K pot <sup>-1</sup> .....	102

## LIST OF FIGURES

Figure	Page
1 The Potassium Cycle in the Soil-Plant System.....	9
2 Tensile Load of Rice Straw Segments (5 cm) from the Internode....	27
3 Temperature Variations of Decomposing Rice Straw and Ambient during Composting.....	33
4 Changes in Organic Matter Remaining and Ash Content of Decomposing Rice Straw during Composting.....	35
5 Changes in C to N Ratio of Decomposing Rice Straw during Composting .....	35
6 Changes in Tensile Load of Indicator Rice Straw during Composting of Rice Straw.....	37
7 Relationships between Organic Matter Remaining and Ash Content with Tensile Load of Indicator Rice Straw during Composting of Rice Straw.....	37
8 Changes in Total and Water Soluble K Contents of Decomposing Rice Straw during Composting.....	39
9 Relationships between Total and Water Soluble K Contents with Tensile Load of Indicator Rice Straw during Composting.....	39
10 Dry Matter Yield of Corn at Different Growth Periods Grown on Soil Treated with Different Rates and Sources of K Fertilizer.....	56
11 The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on Dry Matter Yield of Corn Crop at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	58
12 Crop K Uptake of Corn at Different Growth Periods Grown on Soil Treated with Different Rates and Sources of K Fertilizer.....	59
13 The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on K Uptake by Corn Crop at (a) 14, (b) 28 and (c) 56 After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	60

14	Means for Apparent K Recovery from Muriate of Potash (MOP), Uncomposted (UCRS) and Composted Rice Straw (CRS) for the Corn Crop at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from MOP is 360 mg K pot <sup>-1</sup> .....	63
15	Crop P, Ca and Mg Uptake of Corn at Different Growth Periods Grown on Soil Treated with Different Rates and Sources of K Fertilizer.....	65
16	The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on P Uptake by Corn Crop at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	66
17	The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on Ca Uptake by Corn Crop at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	67
18	The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on Mg Uptake by Corn Crop at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	68
19	Soil Exchangeable and Total K Contents at Different Growth Periods of Corn Crop at Different Rates and Sources of K Fertilizer.....	72
20	The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on Exchangeable K Content of Soil at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	74
21	The Effects of Rate of K Applied from Uncomposted (UCRS) and Composted Rice Straw (CRS) on Total K Content of Soil at (a) 14, (b) 28 and (c) 56 Days After Planting. Standard Rate of K from Muriate of Potash (MOP) is 360 mg K pot <sup>-1</sup> .....	75
22	Changes in Tensile Load of Decomposing Indicator Rice Straw and K Uptake by Corn Crop under Soil Applied with Uncomposted Rice Straw (UCRS) Equivalent to 180, 360 and 540 mg K pot <sup>-1</sup> during the Crop Growth Period of 56 Days.....	78
23	Changes in Tensile Load of Decomposing Indicator Rice Straw and Leaf K Concentration of Corn Crop under Soil Applied with Uncomposted Rice Straw (UCRS) Equivalent to 180, 360 and 540 mg K pot <sup>-1</sup> during the Crop Growth Period of 56 Days.....	81

## LIST OF ABBREVIATIONS

LOI	loss on ignition
TOC	total organic carbon
N	newtons
N	nitrogen
<i>N</i>	Normal
<i>M</i>	Molar
P	phosphorus
K	potassium
Ca	calcium
Mg	magnesium
CEC	cation exchange capacity
BD	bulk density
WHC	water holding capacity
CRS	composted rice straw
UCRS	uncomposted rice straw
MOP	Muriate of Potash
TSP	Triple Super Phosphate
GML	Ground Magnesium Limestone
nm	nano metre
MPa	Mega Pascal
dw	dry weight
dmw	dry matter weight

## CHAPTER I

### INTRODUCTION

Potassium fertilization is important in increasing and sustaining crop production, this is so because the natural K supply of most soils are often unable to meet the K needs of crops for high yield. At practical level, K fertilization can increase the efficient use of other nutrients by crops, particularly of N (Syers, 1998). However in many ASEAN countries, particularly in Indonesia and Malaysia, due to limitation of recoverable K reserves, the availability of inorganic K fertilizers depends heavily on import, the accompanying cost more often than not limit farmers in using K fertilizer. Recoverable K reserves in South and Southeast Asia account for only 7.5 % of the world K reserves (Sheldrick, 1985). Production of manufactured K<sub>2</sub>O fertilizers in 1993 was 0.4 % of total world production and regional imports (excluding Japan) amounted to  $3.6 \times 10^6$  tons K<sub>2</sub>O, or 24 % of the world total (Maene, 1995).

The use of K fertilizer in many ASEAN countries is lower compared to N fertilizer (Christensen, 1995). Recommendations of K addition in most intensively irrigated rice domains are insufficient to replace K removal and as a result there is significant depletion of soil K at many sites (Dobermann *et al.*, 1996). Hence, it is crucial to manage or utilize other K source materials, such as crop residues, that are locally available as an alternative means to reduce the over dependency on inorganic K fertilizer. If properly used, returning crop residues to soil not only can reduce the

use of inorganic (chemical) fertilizers but also has beneficial effect on the improvement of soil physical properties. Balanced fertilization through integrated nutrient supply and management is the most practical and viable technique, which holds the key to sustain yields and crop quality without adversely affecting the environment (Motsara, 1995).

With particular reference to Indonesia, rice (*Oryza sativa* L.) is the most important annual crop, yielding a large amount of rice straw annually. The yield of rice straw (dry basis) ranged from 3.96 to 5.32 tons ha<sup>-1</sup>. The land area of rice based cropping system in 1993, for example, was about 11.01 million ha (Tjiptoningsih and Daulay, 1995). In some croplands, the rice crop is planted in rotation with corn (*Zea mays* L.) with sequence rotation rice-corn (Moentono and Fagi, 1992). At times, a major portion of the rice straw is removed from the field or considered as waste and burnt. Managing rice straw (in term of K source) is essential if sustainability is to be achieved with a small amount of inorganic K fertilizer addition. As an organic material, rice straw naturally has high levels of C and K, besides, other essential plant nutrients. About 80 % K taken up by rice crop is in the straw (De Datta and Gomez, 1982; Dobermann *et al.*, 1996).

Some researches have shown that returning rice straw to soil can contribute considerable K for crops, thus reducing the need of K fertilizer, maintains soil K level and increases the yield of crops (Gill and Sri Adiningsih, 1986; Cox and Uribe, 1992). However, there is lack of information about K supplying characteristic and crop available K from the rice straw. Information of this kind may help in the

understanding of K cycling in soil–crop system. It may also lead to help in increasing efficient rice straw management and K fertilizer utilization.

This thesis reports a research consisting of two experiments on composting of rice straw as a means to promote the straw decomposition and a pot experiment using corn variety PJ-58 as test crop for K uptake planted on the Bungor series soil (Typic Palcudult) in a glasshouse. The objective of the first experiment was to study the change in tensile load of rice straw during rice straw composting, and its possibility to be used as an indicator for the state of decomposition and K released. While the objectives of second experiment were 1) to evaluate the crop available K from composted and uncomposted rice straw (CRS and UCRS) compared to that from Muriate of Potash (MOP) as standard K fertilizer and 2) to evaluate the use of tensile load of rice straw as an indicator to predict K released from rice straw incorporated in the soil, the crop K uptake and leaf K concentration.

## CHAPTER II

### LITERATURE REVIEW

#### Potassium Requirement and the Function in Crop Growth and Yield

Potassium (K) is an essential plant nutrient required in large amount, greater than any other nutrient with except to nitrogen, for metabolism and growth (Tisdale *et al.*, 1993; Plaster, 1997). Potassium has many biochemical and biophysical functions and plays an important part in stimulating photosynthesis; partitioning and storage of assimilate as well as in adaptation of crops to environmental stress. It is the major inorganic constituent in the cell vacuoles contributing to the osmotic and turgor potential of cells. It is an essential cation in all processes that are activated by proton pump (photosynthesis, phloem loading), and also needed for the activation of many enzymes (Beringer, 1982). The functions of K explain the importance of K for crop growth and yield. Sufficiency of K nutrition in crops has beneficial effect on the quality of a wide range of crops (Usherwood, 1985), especially in terms of improved protein quantity and quality; decrease the incidence of plant diseases (Huber and Army, 1985).

Tropical tuber crops such as yam, cassava, sweet potato and coco yam, plantation crops such as sugarcane, coconut, tea, coffee, cocoa, rubber, oil palm and banana, and the cereal crops, particularly paddy, have high K requirements (Sekhon, 1982). In addition, most of vegetable crops such as peas, beans, spinach, sweet corn,



cantaloupe, lettuce, onion, broccoli, brussels sprouts, tomato, carrots and celery also require high K to realize their high yields (Geraldson, 1985). Fast growing, early maturing cultivars have a higher rate requirement than late maturing ones (Halevy, 1976), and improved varieties of crops with greater potential for yield generally require higher supply of nutrients to realize their genetic yield potential (Cook, 1986). The high K requirement of crops, especially during their vegetative growth, demands a sufficient supply of this nutrient to achieve higher yield of the crops (Beringer, 1982).

### **Potassium in Soil**

Potassium is present in igneous, sedimentary and metamorphic rocks; comprises about 2.5 % of the earth's crust in which it is the seventh most abundant element (Sheldrick, 1985). The lithosphere contains an average of 1.9 % K (Rich, 1968). However, the soil content an average of 1.2 % K, being lower than the lithosphere's, due to K lost through weathering. Organic soils are low in K because of their low mineral contents, and average value may be less than 0.03 % K. Young soils, having little weathering, have higher than average K contents (Barber, 1995). The available K in plants is generally only a small fraction of the total available of K in soil. The low fraction of K available is a consequence of both its inaccessibility and its relatively strong bonding in the 2:1 clay mineral structure (Lean and Watson, 1985).